

General

Title

Pressure ulcer: rate per 1,000 discharges.

Source(s)

AHRQ quality indicators. Guide to patient safety indicators [version 3.1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2007 Mar 12. 76 p. (AHRQ Pub; no. 03-R203).

AHRQ quality indicators. Patient safety indicators: technical specifications [version 4.1]. PSI #3 pressure ulcer. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2009 Dec 1. 3 p.

Measure Domain

Primary Measure Domain

Outcome

The validity of measures depends on how they are built. By examining the key building blocks of a measure, you can assess its validity for your purpose. For more information, visit the [Measure Validity](#) page.

Secondary Measure Domain

Does not apply to this measure

Brief Abstract

Description

This measure is used to assess the number of cases of pressure ulcer per 1,000 discharges with a length of stay greater than 4 days.

Rationale

Hospitals in the United States provide the setting for some of life's most pivotal events - the birth of a child, major surgery, treatment for otherwise fatal illnesses. These hospitals house the most sophisticated medical technology in the world and provide state-of-the-art diagnostic and therapeutic

services. But access to these services comes with certain costs. About 30% of personal health care expenditures in the United States go towards hospital care, and the rate of growth in spending for hospital services has only recently leveled out after several years of increases following a half a decade of declining growth. Simultaneously, concerns about the quality of health care services have reached a crescendo with the Institute of Medicine's series of reports describing the problem of medical errors and the need for a complete restructuring of the health care system to improve the quality of care. Policymakers, employers, and consumers have made the quality of care in U.S. hospitals a top priority and have voiced the need to assess, monitor, track, and improve the quality of inpatient care.

Widespread consensus exists that health care organizations can reduce patient injuries by improving the environment for safety from implementing technical changes, such as electronic medical record systems, to improving staff awareness of patient safety risks. Clinical process interventions also have strong evidence for reducing the risk of adverse events related to a patient's exposure to hospital care. Patient Safety Indicators (PSIs), which are based on computerized hospital discharge abstracts from the AHRQ's Healthcare Cost and Utilization Project (HCUP), can be used to better prioritize and evaluate local and national initiatives. Analyses of these and similar inexpensive, readily available administrative data sets may provide a screen for potential medical errors and a method for monitoring trends over time.

The Pressure Ulcer indicator* is intended to flag cases of in-hospital pressure ulcers. Its definition is limited to pressure ulcer as secondary diagnosis to better screen out cases that may be present on admission. In addition, this indicator excludes patients who have a length of stay of 4 days or less, as it is unlikely that a pressure ulcer would develop within this period of time. Finally, this indicator excludes patients who are particularly susceptible to pressure ulcer, namely patients with major skin disorders (Major Diagnostic Category 9 [MDC 9]) and paralysis.

*The following concerns affect the validity of this indicator:

Underreporting or screening: Conditions included in this indicator may not be systematically reported (leading to an artificially low rate) or may be routinely screened for (leading to a higher rate in facilities that screen).

Heterogeneous severity: This indicator includes codes that encompass several levels of severity of a condition that cannot be ascertained by the codes.

Case mix bias: This indicator was felt to be particularly subject to systematic bias, and Diagnosis-Related Group (DRG) and comorbidity risk adjustment may not adequately address the concern.

Refer to the original measure documentation for further information.

Primary Clinical Component

Pressure ulcer

Denominator Description

All medical and surgical discharges, 18 years and older, defined by specific Diagnosis-Related Groups (DRGs) or Medicare Severity DRGs (MS-DRGs)

Exclude cases:

With length of stay of less than 5 days

With principal diagnosis of pressure ulcer or secondary diagnosis present on admission

Major Diagnostic Category (MDC) 9 (Skin, Subcutaneous Tissue, and Breast)

MDC 14 (pregnancy, childbirth, and puerperium)

With any diagnosis of hemiplegia, paraplegia, or quadriplegia

With any diagnosis of spina bifida or anoxic brain damage

With an International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) procedure code for debridement or pedicle graft before or on the same day as the major operating room procedure (surgical cases only)

With any diagnosis of Stage I or Stage II pressure ulcer

Transfer from a hospital (different facility)

Transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF)
Transfer from another health care facility

Note: Refer to the Technical Specifications document for specific ICD-9-CM codes, DRGs and MS-DRGs.

Numerator Description

Discharges with International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) code of pressure ulcer in any secondary diagnosis field among cases meeting the inclusion and exclusion rules for the denominator

Note: Refer to the Technical Specifications document for specific ICD-9-CM codes.

Evidence Supporting the Measure

Evidence Supporting the Criterion of Quality

A formal consensus procedure involving experts in relevant clinical, methodological, and organizational sciences

One or more research studies published in a National Library of Medicine (NLM) indexed, peer-reviewed journal

Evidence Supporting Need for the Measure

Need for the Measure

Variation in quality for the performance measured

Evidence Supporting Need for the Measure

AHRQ quality indicators. Guide to patient safety indicators [version 3.1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2007 Mar 12. 76 p. (AHRQ Pub; no. 03-R203).

State of Use of the Measure

State of Use

Current routine use

Current Use

Internal quality improvement

National reporting

Quality of care research

Application of Measure in its Current Use

Care Setting

Hospitals

Professionals Responsible for Health Care

Nurses

Lowest Level of Health Care Delivery Addressed

Individual Clinicians

Target Population Age

Age greater than or equal to 18 years

Target Population Gender

Either male or female

Stratification by Vulnerable Populations

Unspecified

Characteristics of the Primary Clinical Component

Incidence/Prevalence

Unspecified

Association with Vulnerable Populations

Unspecified

Burden of Illness

Unspecified

Utilization

Unspecified

Costs

Unspecified

Institute of Medicine (IOM) Healthcare Quality Report Categories

IOM Care Need

Getting Better

IOM Domain

Safety

Data Collection for the Measure

Case Finding

Users of care only

Description of Case Finding

All medical and surgical discharges, 18 years and older, defined by specific Diagnosis-Related Groups (DRGs) or Medicare Severity DRGs (MS-DRGs) (see the "Denominator Inclusions/Exclusions" field)

Denominator Sampling Frame

Patients associated with provider

Denominator Inclusions/Exclusions

Inclusions

All medical and surgical discharges, 18 years and older, defined by specific Diagnosis-Related Groups (DRGs) or Medicare Severity DRGs (MS-DRGs)

Exclusions

Exclude cases:

- With length of stay of less than 5 days

- With principal diagnosis of pressure ulcer or secondary diagnosis present on admission

- Major Diagnostic Category (MDC) 9 (Skin, Subcutaneous Tissue, and Breast)

- MDC 14 (pregnancy, childbirth, and puerperium)

- With any diagnosis of hemiplegia, paraplegia, or quadriplegia

- With any diagnosis of spina bifida or anoxic brain damage

- With an International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) procedure code for debridement or pedicle graft before or on the same day as the major operating room procedure (surgical cases only)

- With any diagnosis of Stage I or Stage II pressure ulcer

- Transfer from a hospital (different facility)

- Transfer from a Skilled Nursing Facility (SNF) or Intermediate Care Facility (ICF)

Transfer from another health care facility

Note: Refer to the Technical Specifications document for specific ICD-9-CM codes, DRGs and MS-DRGs.

Relationship of Denominator to Numerator

All cases in the denominator are equally eligible to appear in the numerator

Denominator (Index) Event

Clinical Condition

Institutionalization

Therapeutic Intervention

Denominator Time Window

Time window is a single point in time

Numerator Inclusions/Exclusions

Inclusions

Discharges with International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) code of pressure ulcer in any secondary diagnosis field among cases meeting the inclusion and exclusion rules for the denominator

Note: Refer to the Technical Specifications document for specific ICD-9-CM codes.

Exclusions

Unspecified

Measure Results Under Control of Health Care Professionals, Organizations and/or Policymakers

The measure results are somewhat or substantially under the control of the health care professionals, organizations and/or policymakers to whom the measure applies.

Numerator Time Window

Institutionalization

Data Source

Administrative data

Level of Determination of Quality

Not Individual Case

Outcome Type

Adverse Outcome

Pre-existing Instrument Used

Unspecified

Computation of the Measure

Scoring

Rate

Interpretation of Score

Better quality is associated with a lower score

Allowance for Patient Factors

Analysis by high-risk subgroup (stratification on vulnerable populations)

Analysis by subgroup (stratification on patient factors, geographic factors, etc.)

Risk adjustment method widely or commercially available

Description of Allowance for Patient Factors

Risk adjustment of the data is recommended using age, sex, modified Diagnosis-Related Group (DRG), and comorbidity categories.

Application of multivariate signal extraction (MSX) to smooth risk adjusted rates is also recommended.

Standard of Comparison

External comparison at a point in time

External comparison of time trends

Internal time comparison

Evaluation of Measure Properties

Extent of Measure Testing

The Patient Safety Indicators (PSIs) were evaluated by the project team using empirical analyses to explore the frequency and variation of the indicators, the potential bias, based on limited risk adjustment, and the relationship between indicators. The data sources used in the empirical analyses were the 1997 Florida State Inpatient Database (SID) for initial testing and development and the 1997 Healthcare Cost and Utilization Project (HCUP) State Inpatient Database for 19 States for the final empirical analyses.

All potential indicators were examined empirically by developing and conducting statistical tests for precision, bias, and relatedness of indicators. Three different estimates of hospital performance were calculated for each indicator:

The raw indicator rate was calculated using the number of adverse events in the numerator divided by the number of discharges in the population at risk by hospital.

The raw indicator was adjusted to account for differences among hospitals in age, gender, modified Diagnosis-Related Group (DRG), and comorbidities.

Multivariate signal extraction methods were applied to adjust for reliability by estimating the amount of "noise" (i.e., variation due to random error) relative to the amount of "signal" (i.e., systematic variation in hospital performance or reliability) for each indicator.

The project team constructed a set of statistical tests to examine the precision, bias, and relatedness of indicators for all accepted Provider-level Indicators, and precision and bias for all accepted Area-level Indicators. It should be noted that rates based on fewer than 30 cases in the numerator or the denominator are not reported.

The project team conducted a structured review of each indicator to evaluate the face validity (from a clinical perspective) of the indicators. The methodology for the structured review was adapted from the RAND/UCLA Appropriateness Method and consisted of an initial independent assessment of each indicator by clinician panelists using an initial questionnaire, a conference call among all panelists, followed by a final independent assessment by clinician panelists using the same questionnaire. The review sought to establish *consensual validity*, which "extends face validity from one expert to a panel of experts who examine and rate the appropriateness of each item..." The panel process served to refine definitions of some indicators, add new measures, and dismiss indicators with major concerns from further consideration.

Refer to the original measure documentation for additional details.

Evidence for Reliability/Validity Testing

AHRQ quality indicators. Guide to patient safety indicators [version 3.1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2007 Mar 12. 76 p. (AHRQ Pub; no. 03-R203).

Identifying Information

Original Title

Pressure ulcer (PSI 3).

Measure Collection Name

Agency for Healthcare Research and Quality (AHRQ) Quality Indicators

Measure Set Name

Patient Safety Indicators

Submitter

Agency for Healthcare Research and Quality - Federal Government Agency [U.S.]

Developer

Agency for Healthcare Research and Quality - Federal Government Agency [U.S.]

Funding Source(s)

Agency for Healthcare Research and Quality (AHRQ)

Composition of the Group that Developed the Measure

The Agency for Healthcare Research and Quality (AHRQ) Quality Indicators are in the public domain and the specifications come from multiple sources, including the published and unpublished literature, users, researchers, and other organizations. AHRQ as an agency is responsible for the content of the indicators.

Financial Disclosures/Other Potential Conflicts of Interest

None

Included in

National Healthcare Disparities Report (NHDR)

National Healthcare Quality Report (NHQR)

Adaptation

This indicator was originally proposed by Iezzoni and colleagues (1994) as part of Complications Screening Program (CSP) (CSP 6, "cellulitis or decubitus ulcer"). Needleman and Buerhaus (2001) identified decubitus ulcer as an "outcome potentially sensitive to nursing." The American Nurses Association (1999), its State associations, and the California Nursing Outcomes Coalition have identified the total prevalence of inpatients with Stage I, II, III, or IV pressure ulcers as a "nursing-sensitive quality indicator for acute care settings."

Release Date

2003 Mar

Revision Date

2009 Dec

Measure Status

This is the current release of the measure.

This measure updates previous versions:

AHRQ quality indicators. Guide to patient safety indicators [version 3.0a]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2006 May 1. 78 p. (AHRQ Pub; no. 03-R203).

AHRQ quality indicators. Patient safety indicators: technical specifications [version 3.2]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2008 Mar 10. 107 p.

Source(s)

AHRQ quality indicators. Guide to patient safety indicators [version 3.1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2007 Mar 12. 76 p. (AHRQ Pub; no. 03-R203).

AHRQ quality indicators. Patient safety indicators: technical specifications [version 4.1]. PSI #3 pressure ulcer. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2009 Dec 1. 3 p.

Measure Availability

The individual measure, "Pressure Ulcer (PSI 3)," is published in the "AHRQ Quality Indicators. Guide to Patient Safety Indicators" and "AHRQ Quality Indicators. Patient Safety Indicators: Technical Specifications." These documents are available in Portable Document Format (PDF) from the [Patient Safety Indicators Download](#) page at the Agency for Healthcare Research and Quality (AHRQ) Quality Indicators Web site.

For more information, please contact the QI Support Team at support@qualityindicators.ahrq.gov.

Companion Documents

The following are available:

AHRQ quality indicators. Patient safety indicators: software documentation, SAS [version 4.1]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2009 Dec 2. 37 p. This document is available in Portable Document Format (PDF) from the [Agency for Healthcare Research and Quality \(AHRQ\) Quality Indicators Web site](#) .

Agency for Healthcare Research and Quality SAS® documentation addendum [version 4.1a]. Revisions to AHRQ QI documentation. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2010 Jul 13. 2 p. This document is available in PDF from the [AHRQ Quality Indicators Web site](#) .

AHRQ quality indicators. Software documentation: Windows [version 4.1a]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2010 Jul 2. 97 p. This document is available in PDF from the [AHRQ Quality Indicators Web site](#) .

AHRQ quality indicators. Patient safety quality indicators composite measure workgroup. Final report. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2008 Mar. various p. This document is available in PDF from the [AHRQ Quality Indicators Web site](#) .

AHRQ quality indicators (AHRQ QI). Guidance on using the AHRQ QI for hospital-level comparative reporting [version 1.0]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2009 Jun 30. 41 p. This document is available in PDF from the [AHRQ Quality Indicators Web site](#) .

UCSF-Stanford Evidence-based Practice Center. Davies GM, Geppert J, McClellan M, et al. Refinement of the HCUP quality indicators. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); 2001 May. (Technical review; no. 4). This document is available in PDF from the [AHRQ Quality Indicators Web site](#) .

HCUPnet: a tool for identifying, tracking, and analyzing national hospital statistics. [Web site]. Rockville (MD): Agency for Healthcare Research and Quality (AHRQ); [accessed 2010 Jan 4]. HCUPnet is available from the [AHRQ Web site](#) . See the related [QualityTools](#) summary.

NQMC Status

This NQMC summary was completed by ECRI on October 1, 2003. The information was verified by the

measure developer on October 29, 2003. This NQMC summary was updated by ECRI on February 7, 2005 and on April 11, 2006. The information was verified by the measure developer on July 31, 2006. This NQMC summary was updated by ECRI Institute on June 12, 2007, October 15, 2008 and again on June 21, 2010.

Copyright Statement

No copyright restrictions apply.

Disclaimer

NQMC Disclaimer

The National Quality Measures Clearinghouse[®] (NQMC) does not develop, produce, approve, or endorse the measures represented on this site.

All measures summarized by NQMC and hosted on our site are produced under the auspices of medical specialty societies, relevant professional associations, public and private organizations, other government agencies, health care organizations or plans, individuals, and similar entities.

Measures represented on the NQMC Web site are submitted by measure developers, and are screened solely to determine that they meet the [NQMC Inclusion Criteria](#).

NQMC, AHRQ, and its contractor ECRI Institute make no warranties concerning the content or its reliability and/or validity of the quality measures and related materials represented on this site. Moreover, the views and opinions of developers or authors of measures represented on this site do not necessarily state or reflect those of NQMC, AHRQ, or its contractor, ECRI Institute, and inclusion or hosting of measures in NQMC may not be used for advertising or commercial endorsement purposes.

Readers with questions regarding measure content are directed to contact the measure developer.